

A note on incompressible ionized fluid mixtures

TOMÁŠ ROUBÍČEK

Mathematical Institute, Charles University,
Sokolovská 83, CZ-186 75 Praha 8

and

Institute of Information Theory and Automation, Academy of Sciences,
Pod vodárenskou věží 4, CZ-182 08 Praha 8,

CZECH REPUBLIC

tomas.roubicek@mff.cuni.cz, <http://www.karlin.mff.cuni.cz/~roubicek>

Abstract: The model combining non-Newtonian generalization of the Navier-Stokes equation for barycentric velocity with Nernst-Planck equation for concentrations of particular mutually reacting ionic constituents, the heat equation, and also the Poisson equation for self-induced quasistatic electric field is presented. Existence of weak solutions is outlined and, in a special isothermal case, also uniqueness is proved.

Key Words: non-Newtonian fluids, mixtures, Prigogine description, existence, uniqueness.

1 Introduction

Chemically reacting mixtures represent a framework for modelling of various processes in biology and chemistry. My research in this area has been initiated by J. Nečas who, during many years before he passed away, spoke about “living fluids”, although he never elaborated any concept of such fluids. To compromise thermodynamic amenability and mathematical rigor, the model proposed in [17, 18] uses incompressible Newtonian framework with the barycentric impulse balance, also called Eckart-Prigogine’s [6, 14] concept; in the compressible case, see also [1, 3, 4, 8]. The incompressibility refers here both to each particular constituent and, through volume-additivity hypothesis as in e.g. [11, 16], also to the overall mixture. To cover biological applications on a (sub-) cellular level where intensity of electric field on cell membranes is very high (about 10^7Vm^{-1}), the self-induced electrostatic field must be considered. In comparison with [17, 18] or [19, Sect. 12.6], we consider here a non-Newtonian concept and prove existence of solution for the full system.

2 The model

We consider a mixture of L mutually reacting chemical ionic constituents. Our model consists in a system

of $n+L+2$ differential equations combining the *non-Newtonian* modification of the *Navier-Stokes equation* (balancing the barycentric momentum ρv), the *Nernst-Planck equation* modified for moving media (balancing the mass of particular constituents), the *heat equation* (balancing the internal energy $c_v \theta$), and the quasistatic *Poisson equation* for the electrostatic field (balancing the electric induction $\epsilon \nabla \phi$):

$$\rho \frac{\partial v}{\partial t} + \rho(v \cdot \nabla)v - \text{div} \tau(Dv) + \nabla \pi = -q \nabla \phi, \quad \text{div}(v) = 0, \quad (1a)$$

$$\frac{\partial c_\ell}{\partial t} - \text{div}(d \nabla c_\ell + m c_\ell (e_\ell - q) \nabla \phi - c_\ell v) = r_\ell(c_1, \dots, c_L, \theta), \quad \ell = 1, \dots, L, \quad (1b)$$

$$c_v \frac{\partial \theta}{\partial t} - \text{div}(\kappa \nabla \theta - c_v v \theta) = \tau(Dv) : Dv + d \nabla q \cdot \nabla \phi + \sum_{\ell=1}^L m c_\ell e_\ell^2 |\nabla \phi|^2 - m q^2 |\nabla \phi|^2 - \sum_{\ell=1}^L h_\ell(\theta) r_\ell(c, \theta), \quad (1c)$$

$$\text{div}(\epsilon \nabla \phi) + q = 0, \quad q = \sum_{\ell=1}^L e_\ell c_\ell. \quad (1d)$$

The variables v , π , c_ℓ , θ , ϕ and q have the following meaning:

v barycenter velocity,
 π pressure,
 c_ℓ concentration of ℓ -constituent,
 ϕ electrostatic potential,
 θ temperature,
 q the total electric charge,

where the concentrations c_ℓ are to satisfy

$$\sum_{\ell=1}^L c_\ell = 1, \quad c_\ell \geq 0. \quad (2)$$

In (1c) and later on, c abbreviates (c_1, \dots, c_L) . The meaning of the data is:

$\tau = \tau(Dv)$ stress tensor, $Dv = \frac{1}{2}(\nabla v)^\top + \frac{1}{2}\nabla v$,
 $\rho > 0$ mass density,
 e_ℓ valence (=charge) of ℓ -constituent,
 $\varepsilon > 0$ permittivity,
 $r_\ell(c_1, \dots, c_L, \theta)$ ℓ -constituent production rate,
 $h_\ell = h_\ell(\theta)$ enthalpy of the ℓ -constituent,
 $\kappa > 0$ thermal conductivity,
 $c_v > 0$ heat capacity,
 $d > 0$ a diffusion coefficient, and
 $m > 0$ a mobility coefficient.

The system (1) is to be completed by the initial conditions

$$v(0, \cdot) = v_0, \quad c_\ell(0, \cdot) = c_{\ell 0}, \quad \theta(0, \cdot) = \theta_0 \quad (3)$$

on the considered fixed bounded Lipschitz domain $\Omega \subset \mathbb{R}^n$, and by the boundary conditions corresponding, e.g., to a closed container, which, in some simplified version, leads to:

$$v = 0, \quad c_\ell = c_{\ell\Sigma}, \quad \varepsilon \frac{\partial \phi}{\partial \vec{n}} = \alpha(\phi_\Sigma - \phi), \quad \kappa \frac{\partial \theta}{\partial \vec{n}} = 0 \quad (4)$$

on $\Sigma := (0, T) \times \partial\Omega$, where \vec{n} is the unit outward normal to the boundary $\partial\Omega$ and $c_{\ell\Sigma}$ and ϕ_Σ are prescribed.

3 Remarks to the model

The phenomenological flux $j_\ell := -d\nabla c_\ell + mc_\ell(q - e_\ell)\nabla\phi$ in (1b) equals to $-m(c_\ell\nabla\mu_\ell - f_R)$ where

$$\mu_\ell = e_\ell\phi + \frac{d}{m}\ln(c_\ell), \quad (5)$$

plays the role of an *electrochemical potential* and where

$$f_R := q\nabla\phi \quad (6)$$

is a “*reaction force*” keeping the natural requirement $\sum_{\ell=1}^L j_\ell = 0$ satisfied, which eventually fixes also the

equality constraint in (2). The meaning of the heat sources on the right-hand side of (1c) is: The first term $\tau(Dv) : Dv$ represents the heat production rate due to the loss of kinetic energy by viscosity. The second term $d\nabla q \cdot \nabla\phi$ is the power (per unit volume) of the electric current arising by the diffusion flux, which can create local cooling effects. A global cooling effect seems possible via interaction with the environment if $\alpha \neq 0$, expectedly related with the so-called *Peltier effect*. If $\alpha = 0$, one can however see that the overall production due to this term over Ω is nonnegative: indeed, by using Green’s formula twice, one gets

$$\int_\Omega \nabla q \cdot \nabla\phi \, dx = - \int_\Omega \varepsilon \nabla(\Delta\phi) \cdot \nabla\phi \, dx = \int_\Omega \varepsilon |\Delta\phi|^2 \, dx - \int_\Gamma \varepsilon \Delta\phi \frac{\partial \phi}{\partial \vec{n}} \, dS \geq \int_\Gamma q \alpha (\phi_\Gamma - \phi) \, dS = 0. \quad (7)$$

The third term $\sum_{\ell=1}^L mc_\ell e_\ell^2 |\nabla\phi|^2$ is the power of *Joule’s heat* produced by the electric currents j_ℓ . The fourth term $-mq^2 |\nabla\phi|^2 = -mf_R^2$ is the rate of cooling by the force which balances the volume-additivity constraint, and its influence is presumably very small as usually $|q| \ll \max_{\ell=1, \dots, L} |e_\ell|$. Besides, Joule’s heat always dominates this cooling effect because $\sum_{\ell=1}^L c_\ell e_\ell^2 \geq (\sum_{\ell=1}^L c_\ell e_\ell)^2$ if (2) holds, cf. [17, Remark 2.2]. The last term $\sum_{\ell=1}^L h_\ell(\theta) r_\ell(c, \theta)$ is the heat produced or consumed by chemical reactions.

It should be emphasized that many simplifications are adopted in the presented model: we consider small electrical currents (i.e. magnetic field is neglected), adopt the mentioned volume-additivity and incompressibility assumption, assume the diffusion fluxes independent of other constituent’s gradients (cross-effects are neglected) as well as of the temperature gradient (i.e. Soret’s effect is neglected) and (in agreement with Onsager’s reciprocity principle) also the heat flux independent of the concentration gradients (i.e. Dufour’s effect is neglected), and finally the temperature-independent diffusion coefficients, mobility coefficients, and mass densities that are the same for each constituents, i.e. d , m , and ρ , respectively.

There is a newer and more rational concept by Truesdell [23, 24, 25] balancing impulses $\rho c_\ell v_\ell$ (with v_ℓ denoting the velocity of the ℓ -constituent) of all constituents separately together with interactive forces between them, see also [2, 12, 13, 15, 20, 22]. Then our barycentric velocity v equals to $\sum_{\ell=1}^L c_\ell v_\ell$. Recently, Samohýl [21] derived the model (1) by various simplifications from this Truesdell’s rational model. In particular, [21] showed that the reaction force f_R from (6) in (1b) can be derived from a so-called Hittorf referential system related to the veloc-

ity of a dominant un-charged non-reacting constituent (typically water) after transformation to the barycentric system related to v under the assumptions (among others) of very diluted solution and negligible diffusion velocities.

4 Existence analysis outlined

We naturally assume the mass conservation in all chemical reactions and nonnegative production rate of ℓ th constituent if its concentration vanishes, and the volume-additivity constraint holds for the initial and the boundary conditions, i.e.

$$\sum_{\ell=1}^L r_{\ell}(c_1, \dots, c_L) = 0, \quad (8a)$$

$$r_{\ell}(c_1, \dots, c_{\ell-1}, 0, c_{\ell+1}, \dots, c_L) \geq 0, \quad (8b)$$

$$\sum_{\ell=1}^L c_{\ell 0} = \sum_{\ell=1}^L c_{\ell \Sigma} = 1, \quad c_{\ell 0} \geq 0, \quad c_{\ell \Sigma} \geq 0. \quad (8c)$$

Further, we assume $\tau(D) = \Phi'(|D|^2)$, $\Phi : \mathbb{R} \rightarrow \mathbb{R}^+$, and, for some $\varepsilon > 0$, $C \in \mathbb{R}$, it satisfies

$$\Phi(0) = 0, \quad \Phi'(0) = 0, \quad (9a)$$

$$\Phi''(|D|^2)(B, B) \geq \varepsilon(1 + |D|^{p-2})|B|^2, \quad (9b)$$

$$|\Phi''(|D|^2)| \leq C(1 + |D|^{p-2}) \quad (9c)$$

for any $D, B \in \mathbb{R}^{n \times n}$ symmetric. The Korn inequality and (9a,b) imply (cf. [10, Lemma 2.1]) that, for any $v \in W_0^{1,2}(\Omega; \mathbb{R}^n)$,

$$\int_{\Omega} (\tau(Dv_1) - \tau(Dv_2)) : D(v_1 - v_2) dx \geq \zeta \|\nabla v\|_2^2 \quad (10)$$

for some $\zeta > 0$ depending on ε and on Ω and for $\|\cdot\|_p$ the norm in $L^p(\Omega; \mathbb{R}^{n \times n})$; later, it will also abbreviate the norm in $L^p(\Omega)$ or $L^p(\Omega; \mathbb{R}^n)$.

We will prove the existence of a weak solution by Schauder's fixed point technique like in [17]. We define a retract $K : \mathcal{M} \rightarrow \{\xi \in \mathcal{M}; \xi_{\ell} \geq 0, \ell = 1, \dots, L\}$ by

$$K_{\ell}(\xi) := \frac{\xi_{\ell}^+}{\sum_{k=1}^L \xi_k^+}, \quad \xi_{\ell}^+ := \max(\xi_{\ell}, 0), \quad (11)$$

where \mathcal{M} denotes the affine manifold

$$\mathcal{M} := \left\{ \xi \in \mathbb{R}^L; \sum_{\ell=1}^L \xi_{\ell} = 1 \right\}. \quad (12)$$

Let us note that K is continuous and bounded on \mathcal{M} . Considering $\gamma = (\gamma_1, \dots, \gamma_L) =$ "old" concentrations

and $\vartheta =$ "old" temperature, we define the (v, c, θ, ϕ) as the weak solution to the *de-coupled "retracted" system*:

$$\rho \frac{\partial v}{\partial t} + \rho(v \cdot \nabla)v - \operatorname{div} \tau(Dv) + \nabla \pi = -q \nabla \phi, \quad \operatorname{div}(v) = 0, \quad (13a)$$

$$\frac{\partial c_{\ell}}{\partial t} - \operatorname{div}(d \nabla c_{\ell} + m K_{\ell}(\gamma)(e_{\ell} - q) \nabla \phi - c_{\ell} v) = r_{\ell}(K(\gamma), \vartheta), \quad \ell = 1, \dots, L, \quad (13b)$$

$$c_v \frac{\partial \theta}{\partial t} - \operatorname{div}(\kappa \nabla \theta - c_v v \theta) = \tau(Dv) : Dv + d \sum_{\ell=1}^L e_{\ell} \nabla c_{\ell} \cdot \nabla \phi + m \sum_{\ell=1}^L K_{\ell}(\gamma) e_{\ell}^2 |\nabla \phi|^2 - m q^2 |\nabla \phi|^2 - \sum_{\ell=1}^L h_{\ell}(\vartheta) r_{\ell}(K(\gamma), \vartheta), \quad (13c)$$

$$\operatorname{div}(\varepsilon \nabla \phi) + q = 0, \quad q = \sum_{\ell=1}^L e_{\ell} K_{\ell}(\gamma) \quad (13d)$$

with the initial and boundary conditions (3)–(4). Obviously, given (γ, ϑ) , we are to solve subsequently the (now decoupled) equations (13d), (13a), (13b), and (13c) to obtain ϕ, v, c , and θ , respectively.

As the detailed analysis of the full system (1) is indeed nontrivial and out of the scope of this contribution, we outline it only in a particular case $p = 5/2$. Also, for simplicity we assume

$$r_{\ell}, h_{\ell} \text{ continuous and bounded}, \quad (14)$$

although a sub-linear growth of $r_{\ell}(c, \cdot)$ may be admitted, too; cf. [17]. Let us abbreviate $I := (0, T)$ and $Q = I \times \Omega$.

Proposition 1 *Let the assumptions (8), (9), (14) hold, let $v_0 \in W_{0,\operatorname{div}}^{1,p}(\Omega; \mathbb{R}^n)$, $c_0 \in L^2(\Omega; \mathbb{R}^L)$, $\theta_0 \in L^2(\Omega)$, let Ω be of class C^3 , and α and $\phi_{\Sigma}(t, \cdot)$ be smooth, $n \leq 3$, $p = 5/2$. Let $(\gamma, \vartheta) \in L^2(Q; \mathbb{R}^L) \times L^2(I; W^{1,2}(\Omega))$ be given such that $\sum_{\ell=1}^L \gamma_{\ell} = 1$ a.e. on Q . Then, for some $C < +\infty$ independent of (γ, ϑ) , (13) has a unique weak solution which satisfies*

$$\sigma := \sum_{\ell=1}^L c_{\ell} = 1 \quad \text{a.e. on } Q \quad (15)$$

(although $c_{\ell} \geq 0$ need not hold!), and also the follow-

ing a-priori estimates

$$\|v\|_{L^\infty(I;W^{1,p}(\Omega;\mathbb{R}^n)) \cap L^{\frac{2}{p-1}}(I;W^{2,\frac{6}{p+1}}(\Omega;\mathbb{R}^n))} \leq C \quad (16a)$$

$$\left\| \frac{\partial v}{\partial t} \right\|_{L^2(I;L^2(\Omega;\mathbb{R}^n))} \leq C, \quad (16b)$$

$$\|\theta\|_{L^2(I;W^{1,2}(\Omega)) \cap L^\infty(I;L^2(\Omega))} \leq C, \quad (16c)$$

$$\left\| \frac{\partial \theta}{\partial t} \right\|_{L^2(I;W^{1,2}(\Omega)^*)} \leq C, \quad (16d)$$

$$\|c\|_{L^2(I;W^{1,2}(\Omega;\mathbb{R}^L)) \cap L^\infty(I;L^2(\Omega;\mathbb{R}^L))} \leq C, \quad (16e)$$

$$\left\| \frac{\partial c}{\partial t} \right\|_{L^2(I;W^{1,2}(\Omega;\mathbb{R}^L)^*)} \leq C, \quad (16f)$$

$$\|\phi\|_{L^\infty(I;W^{2,2}(\Omega))} \leq C. \quad (16g)$$

Sketch of the proof. First, we prove (15). By summing (13c) for $\ell = 1, \dots, L$ and by (8a), one gets

$$\frac{\partial \sigma}{\partial t} z - d\Delta \sigma + v \cdot \nabla \sigma = 0 \quad (17)$$

cf. also [17, Formula (3.18)]. Due to (8c), the unique solution to this equation is $\sigma \equiv 1$.

Further, we realize that the charge $q = e \cdot K(\gamma)$ in (13d) is always bounded and, in particular, it is in $L^\infty(I;L^2(\Omega))$, and (16g) follows by usual $W^{2,2}$ -regularity of the Δ -operator with (4). Then also the driving force $q \nabla \phi$ in (13a) is bounded in $L^\infty(I;L^6(\Omega;\mathbb{R}^n))$, hence certainly in $L^2(Q;\mathbb{R}^n)$, and we can use [10] where the estimates (16a,b) have been derived by a very sophisticated usage of a shift technique and a test by a truncated Laplacean.

Testing (13b) by c_ℓ gives (16e) standardly when we realize that the term $\text{div}(mK_\ell(\gamma)(e_\ell - q)\nabla \phi) - r_\ell(K(\gamma), \vartheta)$ is certainly bounded in $L^\infty(I;W^{1,6/5}(\Omega)^*) \subset L^2(I;W^{1,2}(\Omega)^*)$ and when we also use

$$\begin{aligned} \int_\Omega c_\ell v \cdot \nabla c_\ell \, dx &= \frac{1}{2} \int_\Omega v \cdot \nabla c_\ell^2 \, dx \\ &= -\frac{1}{2} \int_\Omega (\text{div } v) c_\ell^2 \, dx = 0. \end{aligned} \quad (18)$$

Then (16f) follows by testing (13b) by arbitrary $z \in L^2(I;W^{1,2}(\Omega))$.

For $p = 5/2$, (16a) is the estimate of ∇v in $L^\infty(I;L^{5/2}(\Omega;\mathbb{R}^n)) \cap L^{4/3}(I;W^{1,12/7}(\Omega;\mathbb{R}^n)) \subset L^\infty(I;L^{5/2}(\Omega;\mathbb{R}^n)) \cap L^{4/3}(I;L^5(\Omega;\mathbb{R}^n))$ which is, by interpolation with the coefficients $(\frac{4}{15}, \frac{11}{15})$, embedded into $L^5(I;L^{3+\varepsilon}(\Omega;\mathbb{R}^n))$, here $\varepsilon = \frac{18}{19}$. Due to (9a,c), $\tau(\text{D}v):\text{D}v$ is then certainly bounded in $L^2(I;L^{6/5}(\Omega))$ which is a subset of the natural "right-hand-side space" $L^2(I;W^{1,2}(\Omega)^*)$ for the heat equation. By

(16e,g), we also know that $(e \cdot \nabla c) \cdot \nabla \phi$ is bounded in $L^2(I;L^{3/2}(\Omega))$. The other three terms on the right-hand side of (13c) are even better. Then (16c) follows standardly by testing (13c) by θ , and (16d) then follows by using a test by arbitrary $z \in L^2(I;W^{1,2}(\Omega))$ for (13c).

Eventually, the uniqueness of solutions to (13b,c,d) follows standardly because these equations are decoupled and linear, while uniqueness for (13a) is non-trivial and has been proved in [10] if $p \geq 9/4$. \square

Proposition 2 *Let the assumptions of Proposition 1 hold, then the mapping $(\gamma, \vartheta) \mapsto (v, c, \theta, \phi)$ with $\sum_{\ell=1}^L \gamma_\ell = 1$ is continuous from the weak topology on \mathcal{W}^{L+1} with*

$$\mathcal{W} := L^2(I;W^{1,2}(\Omega)) \cap W^{1,2}(I;W^{1,2}(\Omega)^*) \quad (19)$$

to the weak topology related to the spaces from the estimates (16).*

Sketch of the proof. Take a sequence $\{(g_k, \vartheta_k)\}_{k \in \mathbb{N}}$ converging weakly to some (γ, ϑ) in \mathcal{W}^{L+1} . Take corresponding $(v_k, c_k, \theta_k, \phi_k)$ and choose a subsequence converging weakly* in the spaces specified in the estimates (16). By Aubin-Lions' compact-embedding theorem, cf. e.g. [19, Lemma 7.7], $\gamma_k \rightarrow \gamma$ strongly in $L^2(I;L^{6-\varepsilon}(\Omega;\mathbb{R}^L))$ with $\varepsilon > 0$, which allows us to pass to the limit $K(\gamma_k) \rightarrow K(\gamma)$ and also ensures $\phi_k \rightarrow \phi$ strongly in $L^{1/\varepsilon}(I;W^{2,2}(\Omega))$. Then we get $|\nabla \phi_k|^2 \rightarrow |\nabla \phi|^2$ in $L^{1/(2\varepsilon)}(I;L^3(\Omega))$ to exploit for (13c). Using again Aubin-Lions' theorem shows $\vartheta_k \rightarrow \vartheta$ strongly in $L^2(I;L^{6-\varepsilon}(\Omega))$, which allows us to pass to the limit $h_\ell(\vartheta_k) \rightarrow h_\ell(\vartheta)$ and $r_\ell(\gamma_k, \vartheta_k) \rightarrow r_\ell(\gamma, \vartheta)$. Moreover, again by Aubin-Lions' theorem and by interpolation like in the proof of Proposition 1, $\nabla v_k \rightarrow \nabla v$ in $L^5(I;L^3(\Omega;\mathbb{R}^n))$ hence $\tau(\text{D}v_k):\text{D}v_k \rightarrow \tau(\text{D}v):\text{D}v$ strongly in $L^2(I;L^{6/5}(\Omega))$, which is essential for the limit passage in (13c). to obtain a conventional weak solution. The limit passage in (13) is then routine. The uniqueness proved in Proposition 1 ensures eventually the convergence of the whole sequence. \square

Proposition 3 *Let again the assumptions of Proposition 1 hold, then the mapping $(\gamma, \vartheta) \mapsto (c, \theta) : C \rightarrow C$ with*

$$\begin{aligned} C := \{ &(c, \theta) \in \mathcal{W}^{L+1}; \|c\|_{\mathcal{W}^L} \leq C, \\ &\|\theta\|_{\mathcal{W}} \leq C, c(\cdot, \cdot) \in \mathcal{M} \text{ a.e. on } Q \} \end{aligned} \quad (20)$$

with C from (16c-f) has a fixed point (c, θ) and, considering the corresponding ϕ and v , the quadruple (v, c, θ, ϕ) is a weak solution to (1)–(4).

Sketch of the proof. We use C equipped with the weak topology. The fixed point then exists by Schauder's theorem (in Tikhonov's modification).

Finally, by testing by c_ℓ^- the resulted equation for c_ℓ , i.e. (13b) with $K(c)$ in place of $K(\gamma)$, we obtain $c_\ell^- = 0$ if (8b,c) is taken into account. Hence (2) is proved, and $c_\ell = K_\ell(c)$, so that the retract K can eventually be forgotten in the this fixed point. \square

5 Uniqueness in the isothermal case

Let us confine ourselves on the case that the temperature variations can be neglected, hence instead of $r_\ell = r_\ell(c, \theta)$ we consider only $r_\ell = r_\ell(c)$; it is certainly well satisfied, e.g., in biological applications on cellular level. Then (1) decouples to (1a,b,d) and (1c). To show uniqueness, it suffices to consider only (1a,b,d) because (1c) will follow.

Proposition 4 *Let (9c) hold, r_ℓ be Lipschitz continuous, Ω be of the C^3 -class, α and ϕ_Σ be smooth, $n \leq 3$, and $p \geq 5/2$. Then there is at most one weak solution to the problem (1a,b,d),(3),(4).*

Proof. For notational simplicity, let $p=1$. Recall that $q = \sum_{\ell=1}^L e_\ell c_\ell =: e \cdot u$. Consider the two weak solutions (ϕ^1, c^1, v^1) and (ϕ^2, c^2, v^2) to (1a,b,d), and denote $\phi^{12} := \phi^1 - \phi^2$, $c^{12} := c^1 - c^2$, and $v^{12} := v^1 - v^2$. Test the difference of (1a) (resp. (1b)) written for two solutions by v^{12} (resp. c_ℓ^{12}), and use (10) to get:

$$\begin{aligned} & \frac{d}{dt} \left(\|v^{12}\|_2^2 + \sum_{\ell=1}^L \|c_\ell^{12}\|_2^2 \right) + \zeta \|\nabla v^{12}\|_2^2 \\ & + d \sum_{\ell=1}^L \|\nabla c_\ell^{12}\|_2^2 = \int_\Omega \left(((v^2 \cdot \nabla)v^2 - (v^1 \cdot \nabla)v^1) \cdot v^{12} \right. \\ & + (q^2 \nabla \phi^2 - q^1 \nabla \phi^1) \cdot v^{12} \\ & + m \sum_{\ell=1}^L \left(c_\ell^2 (e_\ell - q^2) \nabla \phi^2 - c_\ell^1 (e_\ell - q^1) \nabla \phi^1 \right) \cdot \nabla c_\ell^{12} \\ & + \sum_{\ell=1}^L (c_\ell^1 v^1 - c_\ell^2 v^2) \nabla c_\ell^{12} \\ & \left. + (r(c^1) - r(c^2)) \cdot c^{12} \right) dx =: I_1 + \dots + I_5 \quad (21) \end{aligned}$$

The term I_1 in (21), arising from the convective term, can be handled as in [9, Theorem 4.29] modified with zero-Dirichlet boundary condition provided $p \geq 5/2$, namely

$$\begin{aligned} I_1 & = - \int_\Omega (v^{12} \cdot \nabla) v^1 \cdot v^{12} dx \\ & \leq \varepsilon \|\nabla v^{12}\|_2^2 + C_\varepsilon \|v^1\|_p^{2p/(2p-n)} \|\nabla v^{12}\|_2^2 \end{aligned}$$

for $\varepsilon < \zeta$ and then treated by Gronwall's inequality.

Furthermore, from (1d) we get $\phi^{12} = \Delta^{-1}(e \cdot c^{12})$ where Δ^{-1} denotes the inverse operator to Δ under the homogeneous boundary conditions (4), i.e. $\varepsilon \partial \phi / \partial \bar{n} + \alpha \phi = 0$. Estimate the term I_2 in (21), for each $\ell = 1, \dots, L$, as

$$\begin{aligned} I_2 & := \int_\Omega \left(c_\ell^2 \nabla \phi^2 - c_\ell^1 \nabla \phi^1 \right) \cdot v^{12} dx \\ & \leq \frac{1}{4\varepsilon} \|c_\ell^{12}\|_2^2 \|\nabla \phi^1\|_4^2 + \varepsilon \|v^{12}\|_4^2 \\ & \quad + \|c_\ell^2\|_\infty^2 \|\nabla \Delta^{-1}(e \cdot c^{12})\|_2^2 + \frac{1}{4} \|v^{12}\|_2^2 =: T_1 + \dots T_4. \end{aligned}$$

By (16e), $q \in L^2(I; W^{1,2}(\Omega))$, and then $\nabla \phi^1 \in L^2(I; W^{2,2}(\Omega; \mathbb{R}^n)) \subset L^2(I; L^4(\Omega; \mathbb{R}^n))$ for $n \leq 3$ (here even $n \leq 8$ is allowed) through standard $W^{3,2}$ -regularity results for the linear boundary-value problem (1d)-(4). Then the term T_1 will be handled by Gronwall's inequality. As to $T_2 \leq \varepsilon N^2 \|\nabla v^{12}\|_2^2$, we will absorb it in the respective term coming from the viscosity term (1a) if $\varepsilon < \zeta/N^2$ where N is the norm of the embedding $W^{1,2}(\Omega) \subset L^4(\Omega)$. As to T_3 , we use $\|\nabla \phi^{12}\|_2 \leq C \|e \cdot c^{12}\|_2$ with some C depending on Ω and on α , and then will handle it together with T_4 by Gronwall's inequality. Now we estimate the terms $I_{3\ell}$ with $I_3 = \sum_{\ell=1}^L I_{3\ell}$ in (21) as

$$\begin{aligned} \frac{I_{3\ell}}{m} & := \int_\Omega \left(c_\ell^1 (e_\ell - q^1) \nabla \phi^1 - c_\ell^2 (e_\ell - q^2) \nabla \phi^2 \right) \cdot \nabla c_\ell^{12} dx \\ & \leq \frac{3m}{d} \|c_\ell^{12}\|_2^2 \|e_\ell - q^1\|_\infty^2 \|\nabla \phi^1\|_\infty^2 \\ & \quad + \frac{3m}{d} \|c_\ell^2\|_\infty^2 \|e \cdot c^{12}\|_2^2 \|\nabla \phi^1\|_\infty^2 \\ & \quad + \frac{3m}{d} \|c_\ell^2\|_\infty^2 \|e_\ell - q^2\|_\infty^2 \|\nabla \phi^{12}\|_2^2 \\ & \quad + \frac{d}{4m} \|\nabla c_\ell^{12}\|_2^2 = T_1 + \dots + T_4. \quad (22) \end{aligned}$$

Now we employ the regularity of $\Delta^{-1} : L^\infty(\Omega) \rightarrow W^{1,\infty}(\Omega)$; this follows by the standard $W^{2,p}$ -regularity theory with $p > n$, cf. e.g. [7], so that $\nabla \phi^1 \in L^\infty(Q; \mathbb{R}^n)$, which is needed for both T_1 and T_2 . These terms are then to be treated by Gronwall's inequality. As to T_3 , estimate $\|\nabla \phi^{12}\|_2^2 \leq C \|e \cdot c^{12}\|_2^2$, which will lead to Gronwall's inequality, while T_4 is to be absorbed in the left-hand side. Further, using also (18) (here with v^1 and c_ℓ^{12} instead of v and c_ℓ , respectively) we estimate $I_{4\ell}$ in the term $I_4 = \sum_{\ell=1}^L I_{4\ell}$ in (21) as

$$\begin{aligned} I_{4\ell} & := \int_\Omega (c_\ell^1 v^1 - c_\ell^2 v^2) \cdot \nabla c_\ell^{12} dx = \int_\Omega c_\ell^2 v^{12} \cdot \nabla c_\ell^{12} dx \\ & \leq \frac{1}{d} \|c_\ell^2\|_\infty^2 \|v^{12}\|_2^2 + \frac{d}{4} \|\nabla c_\ell^{12}\|_2^2. \quad (23) \end{aligned}$$

Eventually, denoting by L_r the Lipschitz constant of $r : \mathbb{R}^L \rightarrow \mathbb{R}^L$, we estimate the term I_5 in as

$$I_5 := \int_{\Omega} (r(c^1) - r(c^2)) \cdot c^{12} dx \leq L_r \|c^{12}\|_2^2. \quad (24)$$

Then we sum $I_1 + \dots + I_5$ and use the mentioned Gronwall's inequality to obtain both $v^{12} = 0$ and $c_\ell^{12} = 0$. \square

In fact, more sophisticated technique from [10] for I_1 allows even for $p \geq \frac{9}{4}$. For $p = \frac{9}{4}$, the existence seems to hold, too; ε in the proof of Proposition 1 is then 0.0738. Yet, e.g., $p > 3$ does not seem to work. In the isothermal case, the existence of a weak solution was shown also in [18] or [19, Sect.12.6] in the Navier-Stokes case (i.e. $p = 2$ was admitted).

Acknowledgments: This work has been partly supported by the grants 201/03/0934 (GA ČR) and MSM 0021620839 (MŠMT ČR).

References:

- [1] L. Andrej, I. Dvořák, F. Maršík: *Biotermodynamika*. Akademia, Praha, 1982. (Revised edition: F.Maršík, I.Dvořák, Akademia, Praha, 1998).
- [2] R.J. Atkin, R.E. Craine: Continuum theories of mixtures: basic theory and historical development. *Q. J. Mech. Appl. Math.* **29** (1976), 209-244.
- [3] R. Balescu: *Equilibrium and Nonequilibrium Statistical Mechanics*. Wiley, New York, 1975.
- [4] R.S. deGroot, P. Mazur: *Non-equilibrium Thermodynamics*. North-Holland, Amsterdam, 1962.
- [5] D.S. Drumheller, A. Bedford: A thermomechanical theory for reacting immiscible mixtures. *Archive Rat. Mech. Anal.* **73** (1980), 257-284.
- [6] C.Eckart: The thermodynamics of irreversible processes. II.Fluid mixtures. *Physical Rev.* **58** (1940), 269-275.
- [7] D.Gilbarg, N.S.Trudinger: *Elliptic Partial differential Equations of Second Order*. Springer, Berlin, 2nd ed., 1983; revised printing 2001.
- [8] V. Giovangigli: *Multicomponent Flow Modeling*. Birkhäuser, Boston, 1999.
- [9] J. Málek, J. Nečas, M. Rokyta, M. Růžička: *Weak and measure-valued solutions to evolution partial differential equations*. Chapman & Hall, London, 1996.
- [10] J. Málek, J. Nečas, M. Růžička: On weak solutions to a class of non-Newtonian incompressible fluids in bounded three-dimensional domains: the case $p \geq 2$. *Adv. in Diff. Equations* **6** (2001), 257-302.
- [11] N. Mills: Incompressible mixtures of Newtonian fluids. *Int. J. Engng. Sci.* **4** (1966), 97-112.
- [12] I. Müller: A thermodynamical theory of mixtures of fluids. *Archive Rat. Mech. Anal.* **28** (1967), 1-39.
- [13] I. Müller, T. Ruggeri: *Rational Extended Thermodynamics*. (2nd ed.) Springer, New York, 1998.
- [14] I. Prigogine: *Étude Thermodynamique des Processes Irreversibles*. Desoer, Lieg, 1947.
- [15] K.R. Rajagopal, L. Tao: *Mechanics of Mixtures*. World Scientific, River Edge, 1995.
- [16] K.R. Rajagopal, A.S. Wineman, M. Gandhi: On boundary conditions for a certain class of problems in mixture theory. *Int. J. Eng. Sci.* **24** (1986), 1453-1463.
- [17] T. Roubíček: Incompressible ionized fluid mixtures. *Continuum Mech. Thermodyn.*, submitted.
- [18] T. Roubíček: Incompressible fluid mixtures of ionized constituents. In: *Trends in Applic. of Math. to Mechanics*. (Eds.Y.Wang, K.Hutter), Shaker, Aachen, 2005, pp.429-440.
- [19] T. Roubíček: *Nonlinear partial differential equations with applications*. Birkhäuser, Basel, 2005, in print.
- [20] I. Samohýl: *Thermodynamics of Irreversible Processes in Fluid Mixtures*. Teubner, Leipzig, 1987.
- [21] I. Samohýl: Application of Truesdell's model of mixture to ionic liquid mixture. *Comp. Math Appl.*, submitted.
- [22] I. Samohýl, M. Šilhavý: Mixture invariance and its applications. *Archive Rat. Mech. Anal.* **109** (1990), 299-321.
- [23] C. Truesdell, W. Noll: *The nonlinear field theories of mechanics*. Handbuch der Physik III/3, Springer, Berlin, 1965.
- [24] C. Truesdell, K.R. Rajagopal: *An introduction to the Mechanics of Fluids*. Birkhäuser, Boston, 2000.
- [25] C. Truesdell, R. Toupin: *The Classical Field Theories*. Handbuch der Physik III/1, Springer, Berlin, 1960.